



In-vitro Anti-inflammatory activity of Siddha formulation Pirrandai Chooranam by Protein (albumin) denaturation assay.

*Kasthuri M¹, Venkatesh R¹, Balamurugan A²

¹PG Scholar, Department of Noinaadal, ²Lecturer Grade II, Department of Noinaadal,

Government Siddha Medical College, Palayamkottai, Tirunelveli.

ABSTRACT

Inflammatory response is a protective response to injury, irritant or infection and is required to remove the injurious stimuli as well as initiate the healing process. It is a complex process and it associated with increased vascular permeability, protein denaturation and membrane alteration. The common symptoms of arthritis is pain, which is due to the inflammation of joints line. Pirrandai chooranam is a polyherbal Siddha formulation, indicated in Siddha text Siddha vaithiya thirattu which contain 9 ingredients of plant origin. It was mainly indicated for the treatment of Vatham. The aim of the study is to evaluate the Anti-inflammatory activity of Anti-inflammatory activity by using Protein (Albumin) denaturation technique. The reaction mixture consisted of bovine serum albumin (5% aqueous solution) and test sample Pirrandai Chooranam (PC) at varying concentration ranges from 100 to 500 µg/ml and standard Diclofenac sodium at the concentration of 100 µg /ml. The results of the study showed that the test drug was effective in inhibiting heat induced albumin denaturation. The Concentration range of VKI at 100, 200, 300, 400 and 500 µg/ml produce significant inhibition of protein denaturation in concentration dependent manner. Maximum percentage of protein inhibition about 74.25 ± 2.81 % was observed at 500 µg/ml, when compared to Diclofenac sodium, a standard anti-inflammatory agent with the maximum inhibition about 96.57 ± 1.27 at the concentration of 100 µg/ml. Hence this study result offers the Pirrandai Chooranam (PC) possess significant Anti-inflammatory activity in Protein (Albumin) denaturation technique.

Keywords:

Pirrandai, Anti-inflammatory, Siddha medicine

Address for correspondence:

Kasthuri M
PG Scholar

CODEN : IJRPHR

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: publisher@ijrphr.com

To access this article online

Website : <http://www.ijrphr.com/>

DOI : 10.121/ijrphr/05.0101.543

Quick response code



INTRODUCTION

Inflammation is a body defense reaction in order to eliminate (or) limit the spread of injuries agents, followed by removal of necrosied cells and tissues. The response consists of changes in blood flow, an increase permeability of blood vessels, migration of fluid proteins and leukocytes from the circulation to site of tissue damage. The causative agents of inflammations are, Infective agents like bacteria, virus, fungi, parasites and their toxins; Immunological agents like cell mediated antigen, antibody reaction; physical agents like heat, cold, radiation and mechanical traumas; Chemical agents like organic and inorganic poisons, inert materials such as foreign bodies. Signs of inflammation: There is 5 cardinal sign in inflammation, (i,e) rubor (redness), tumor (swelling), calor (heat), dolor (pain), functio laesa (less of function). Types of inflammation: Depending upon the defense capacity of the host and duration of response, inflammation is classified into Acute and Chronic. Acute inflammation is an early body response to harmful stimuli and is achieved by increased movement of plasma and Leukocytes (especially granulocytes) from the blood into the injured tissue.

Chronic inflammation is a prolonged process in which tissues inflammation and destruction occur at the same time. It is due to recurrent attacks of acute inflammation (or) followed by acute inflammation. The hallmark of chronic inflammation is the infiltration of tissue site by macrophages, lymphocytes and plasma cells. Macrophages are the principle cells involved in chronic inflammation and produce many effects that contribute to the progression of tissue damage and consequent functional damage.

Rheumatoid arthritis is a chronic, progressive inflammatory autoimmune disease. It affects 0.5 to 1% of population all over the world. RA may be more prevalent in Rural population than in Urban area (R Handa et al. 2016). In a review of RA epidemiology worldwide published in 2006, Almanos et al, compared the incidence and prevalence of RA, based on ACR criteria, across four categories, North America, Northern Europe, Southern Europe and developing countries. In India local survey in Delhi shows prevalence of this disease affecting 0.75 % population. The Mean age onset of the disease in Female is 38± 12.4 years and Male is 44.8 ± 13.12years. The onset of

disease is insidious, beginning with prodrome of fatigue, weakness, joint stiffness, vague arthralgia and myalgia. This is followed by pain and swelling of joints of hands, wrist, feet, ankle and elbow joints. The typical presentation of RA is smaller joints swelling especially proximal Interphalangeal and Metacarpophalangeal joints are affected more severely. The persist inflammation leads to erosive joint damage and functional impairments in vast majority of patients. In later stage, RA produces systemic manifestations such as haematologic, pulmonary, neurological and cardiovascular abnormalities. Siddha system of medicine is one of the oldest systems of medicine in South India, especially in Tamil Nadu. The human body is based on the five physical elements i.e. Panja bootham – Earth, Water, Fire, Air and Space and three humors i.e. Vatham, Pitham and Kapham. The causes of all diseases or the pathological conditions of human body are to sought in the abnormalities of these three varieties of materials. Siddhars classified the diseases into 4448 based on Muktutam. Saint Yugi in Yugi Vaithya Sindhamani, classified 80 types of Vatha diseases.

MATERIALS AND METHODS

Purification of raw drugs

The purification of this drug is as follows as illustrated in *Gunapadam Thathu Jeeva Vagupu* and *Gunapadam mooligaiyiyal*.

Source of raw drugs:

The plant of PC was bought from the traditional raw drug store Tirunelveli. The raw drugs taken for study were authenticated by the concerned Botanist of Medicinal Botany Department, Government Siddha Medical College, Palayamkottai.

In-vitro anti-inflammatory activity of PC

In-vitro anti-inflammatory activity PC as studied using albumin denaturation technique. The reaction mixture consisted of bovine serum albumin (5% aqueous solution) and test sample chloroform extract of PC at varying concentration ranges from 100 to 500 µg/ml along with standard Diclofenac sodium at the concentration of 100 µg /ml of final volume. pH was adjusted by using a small amount of 1N Hydrochloric acid. The samples were incubated at 37°C for 20 min and then heated at 57°C for 3 min. After cooling the sample, 2.5 ml of phosphate buffer

solution was added into each test tube. Turbidity developed was measured spectrophotometrically at 660 nm, for control distilled water was used instead of test sample while product control tests lacked bovine serum albumin. The experiment was performed in triplicate.

The Percentage protection from denaturation is calculated by using the formulae

$$\left[\frac{(A)_{\text{control}} - (A)_{\text{sample}}}{(A)_{\text{control}}} \right] \times 100.$$

Table. 1. List of Ingredients of Pirandai chooranam

S.No	Tamil name	Part used	Family	Chemical name / Botanical name / Zoological name	Quantity
1	Pirrandai	Stem	Vitaceae	Cissus quadrangularis	9 part
2	Chukku	Rhizome	Zingiberaceae	Zingiber officinale	1 part
3	Pepper	Seed	Piperaceae	Piper nigrum	1 part
4	Thippili	Fruit	Piperaceae	Piper longum	1 part
5	Arathai	Rhizome	Zingiberaceae	Alpinia galanga	1 part
6	Omam	Seed	Apiaceae	Trachyspermum roxburghianum	1 part
7	Ilavangam	Flower	Myrtaceae	Syzygium aromaticum	1 part
8	Kothamalli	Seed	Apiaceae	Coriandrum sativum	1 part
9	Kunkuma Poo	Flower	Iridaceae	Crocus sativus	1 part
10	Induppu	Rock Salt	--	Sodium chloride impura	1 part

Statistical analysis

Results are expressed as Mean \pm SD. The difference between experimental groups was compared by One-Way Analysis of Variance (ANOVA) followed by Dunnet Multiple comparison test.

RESULTS AND DISCUSSION

Protein denaturation is a process in which proteins lose their tertiary structure and secondary structure by application of external stress or compound, such as strong acid or base, a concentrated inorganic salt, an organic solvent or heat. Most biological proteins lose their biological function when denatured. Denaturation of proteins is a well-documented cause of inflammation. According to Opie et al 1962, tissue injury during life might be preferable to denaturation of the protein constituents of cell and intracellular substance. Mechanism of denaturation is involved in alteration of electrostatic force, hydrogen, hydrophobic and di-sulphide bonds.

Some inflammatory diseases and arthritis like Rheumatoid arthritis, denaturation of protein in tissues may be the causes of production of auto-antigens. So the protein denaturation is the marker for the inflammation and arthritis. The ability of a substance to inhibit the protein

denaturation, that substance may be possible to prevent the inflammation (i.e. anti-inflammatory substance).

The Concentration range of PC at 100, 200, 300, 400 and 500 $\mu\text{g/ml}$ produce significant inhibition of protein denaturation in concentration dependent manner. The inhibitory effect of different concentration of PC on protein denaturation as shown in Table 2. Maximum percentage inhibition of about $74.25 \pm 2.817\%$ was observed at 500 $\mu\text{g/ml}$ when compare to that of the Diclofenac sodium, a standard anti-inflammatory agent with the maximum inhibition $96.57 \pm 1.27\%$ at the concentration of 100 $\mu\text{g/ml}$. The result obtained from the study clearly indicates that the test drug PC was effective in inhibiting heat induced albumin denaturation

From the result of the study, it was concluded that the test drug PC possess significant anti-inflammatory property in dose dependent manner.

Figure 1. Percentage Inhibition of Protein Denaturation by PC and Standard




Table: 2 Inhibitory effect of different concentration of PC and Diclofenac sodium on protein denaturation

Concentration in µg/ml	Percentage Inhibition of Protein Denaturation
PC 100	12.62 ± 2.56
PC 200	23.31 ± 1.76
PC 300	34.81 ± 1.44
PC 400	54.16 ± 1.61
PC 500	74.25 ± 2.81
Diclofenac sodium (100 µg)	96.57 ± 1.27

Each value represents the mean ± SD. N=3

The Anti – inflammatory activity of VKI is may be due to the presence of active principles, such as Polyphenolic compounds, Alkaloids, Tannins and Flavonoids. Hence, it can be used for management of inflammatory diseases. Inflammation is a part of healing process. However, sometimes inflammation persist for longer time, it produce harmful effect than benefit. Chronic inflammation increase the risk of several serious diseases, such as Rheumatoid arthritis, Asthma, Coronary heart disease, Cancer etc..

There are many drugs used for the treatment of inflammation. NSAID (such as Ibuprofen, Diclofenac, Indomethacin, Aspirin) are commonly used for the management of inflammation. They inhibit the Cyclo-Oxygenase (COX) and Prostaglandin H synthase enzyme,

which convert arachidonic acid, derived from membrane phospholipids, to prostaglandin and leucotriens by COX and 5- lipoxygenase pathway respectively. These are very effective for pain and inflammation, they are not thought to have a disease modifying effect in but chronic uses of these drugs produce some adverse effect. The adverse effects are most commonly happening, after the long dose over a long period of usage and/or without prescriptions of physician

CONCLUSION

In Siddha literature, no of drugs (originated from plants, animals, minerals and metals) are available for chronic inflammation. Among these, plant originated drugs are easily available and low expenses. PC is one of the plant originated formulation. All the ingredients of this formulation chiefly indicated for Vatha diseases in Siddha text Gunapadam Mooligai Vaguppu. The herbal drugs also had proven by its Analgesic and Anti – Inflammatory activities. Now the present study, the formulation PC was scientifically proved for its anti- inflammatory activity.

ACKNOWLEDGEMENT

I wish to thanks to Faculty in Department of *Noinaadal*, Government Siddha Medical College, Palayamkottai..

CONFLICT OF INTEREST

None Declared

REFERENCES

1. Regan L. Bailey et al., The epidemiology of Global Micronutrient Deficiencies; National Institute of Health (NIH). *Annals of nutrition and metabolism* 2015; 66 (suppl 2); 22-33. <http://doi.org/10.1159/000371618>.
2. Nguyen PH, Scott S, Avula R, et al., Trends and drivers of change in the prevalence of anaemia among 1 million women and children in india,2006 to 2016. *BMJ Global health* September 2018; 3: page 1- 12.
3. Dr. Ann B Bruner et al., Randomized study of cognitive effects of Iron supplementation in Non – Anaemic Iron – deficient adolescent girls, *THE LANCET*; Vol 348, Issue 9033, October 12, 1996: pages 992 – 96. DOI: [https://doi.org/10.1016/S0140-6736\(96\)02341-0](https://doi.org/10.1016/S0140-6736(96)02341-0).
4. Himanisharma et al., Major Correlates of Anaemia among Women (age 15 – 49) in India and spatial variation, evidence from national family health survey

- 4; journal of women's health care 2018, 7; 4;
<https://www.researchgate.net/publication/327842055>.
5. The Global Prevalence of Anaemia in 2011, WHO Document production services, Geneva: Switzerland. World health organization 2015.
 6. S. P. Ramachandran, *Paandu roganithanam, Yugi vaithiya chinthamani (Moolamumuraium)*; Thamarainoolagam, 2nd edition, July 2013; page no: 201.
 7. Abhimanyukumar, Asishkumargarai, A clinical study on *Paandu roga*, Iron deficiency anemia with Trikatrayadilauha suspension in children; Journal of Ayurveda & integrative medicine. 2012 Oct – Dec; 3 (4): 215 – 222.
 8. J. Seetharamprasath, *roganithanam, paandu, Anubava vaithiya deva ragasiyam* four parts, B. Rathinanaayakkar and sons; first edition 2014; part III, Page no: 436.
 9. C. Kannusaampillai, *Sarakkusuthimuraigal, Chikitcharathnadeepamennum vaithiya nool*, B. Rathnanaayakkar and sons, first edition reprinted 2014, page no: 30.